

University of Groningen

Multiplicity of 3-Ketosteroid-9 alpha-Hydroxylase Enzymes in *Rhodococcus rhodochrous* DSM43269 for Specific Degradation of Different Classes of Steroids

Petrusma, Mirjan; Hessels, Gerda; Dijkhuizen, Lubbert; van der Geize, Robert

Published in:
Journal of Bacteriology

DOI:
[10.1128/JB.00274-11](https://doi.org/10.1128/JB.00274-11)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2011

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Petrusma, M., Hessels, G., Dijkhuizen, L., & van der Geize, R. (2011). Multiplicity of 3-Ketosteroid-9 alpha-Hydroxylase Enzymes in *Rhodococcus rhodochrous* DSM43269 for Specific Degradation of Different Classes of Steroids. *Journal of Bacteriology*, 193(15), 3931-3940. <https://doi.org/10.1128/JB.00274-11>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

Supplemental Table 1. Plasmids used in this study.

Construct	Description	Reference/origin
pRRE1	<i>E. coli</i> - <i>Rhodococcus</i> shuttle vector	Rosłonec <i>et al.</i> (2009) (26)
pKSH602	Clone isolated from genomic library of DSM43269 carrying <i>kshA1</i>	Wilbrink <i>et al.</i> , submitted
pA1rho9	pRRE1 carrying DSM43269 genomic region of <i>kshA1</i> for functional complementation of RG32; 6,296 bp <i>EcoRI</i> fragment of pKSH602 ligated in pRRE1 <i>EcoRI</i>	This study
pKSH509	Clone isolated from genomic library of DSM43269 carrying <i>kshA2</i>	Wilbrink <i>et al.</i> , submitted
pA2rho5	pRRE1 carrying DSM43269 genomic region of <i>kshA2</i> for functional complementation of RG32; 5,291 bp <i>EcoRV/HindIII</i> fragment of pKSH509 ligated in pRRE1 <i>EcoRV/HindIII</i>	This study
pKSH800	Clone isolated from genomic library of DSM43269 carrying <i>kshA3</i>	Wilbrink <i>et al.</i> , submitted
pZero2.1	General cloning vector	Invitrogen
pKSH801	DSM43269 genomic region of <i>kshA3</i> ; \pm 3 kb <i>BamHI</i> fragment of pKSH800 ligated in pZero <i>BamHI</i>	This study
pA3rho5	pRRE1 derivative carrying DSM43269 genomic region of <i>kshA3</i> for functional complementation of RG32; 2,401 bp <i>EcoRV</i> fragment of pKSH801 ligated in pRRE1 <i>EcoRV</i>	This study
pKSH628	Clone from genomic library of DSM43269 carrying <i>kshA4</i>	Wilbrink <i>et al.</i> , submitted
pKSH629	Clone from genomic library of DSM43269 carrying <i>kshA4</i>	Wilbrink <i>et al.</i> , submitted
pA4rho3	DSM43269 genomic region of <i>kshA4</i> ; ligation of 3,182 bp <i>XmnI/DraIII</i> fragment of pKSH628 with 4,320 bp <i>XmnI/DraIII</i> fragment of pKSH629	This study
pA4rho15	pRRE1 carrying DSM43269 genomic region of <i>kshA4</i> for functional complementation RG32; 4,251 bp <i>EcoRI/HindIII</i> fragment of pA4rho3 ligated in pRRE1 <i>EcoRI/HindIII</i>	This study
pBrho5	<i>kshB</i> in pET15b	Petrusma <i>et al.</i> (2009) (24)
pBSKS	pBluescript (II), cloning vector	Stratagene
pKSH814	cloning vector carrying <i>kshA</i> ; KshARho-1-F + -R PCR (Vent polymerase) product (<i>kshA1</i>) ligated in pBSKS <i>EcoRV</i>	This study
pET15b	T7 expression plasmid	Novagen
pKSH816	<i>kshA1</i> expression plasmid; 1,190 bp <i>NdeI/HindIII</i> fragment (<i>kshA1</i>) pKSH 814 ligated in pET15b <i>NdeI/HindIII</i> .	This study
pA1rho3	co-expression construct <i>kshA1-kshB</i> ; 1,289 bp <i>HindIII/XbaI</i> fragment pKsh816 ligated in pBrho5 <i>HindIII/SpeI</i>	This study
pKSH660	cloning vector carrying <i>kshA</i> ; KshARho-2-F2 + -R PCR (High Fidelity polymerase) product (<i>kshA2</i>) ligated in pZero <i>EcoRV</i>	This study
pKSH661	<i>kshA2</i> expression plasmid; 1,268 bp <i>NdeI/HindIII</i> fragment of pKSH 660 ligated in pET15b <i>NdeI/HindIII</i>	This study
pA2rho2	co-expression construct <i>kshA2-kshB</i> ; 1,216 bp <i>HindIII/XbaI</i> fragment of pBrho5 ligated in pKsh661 <i>HindIII/SpeI</i>	This study
pCR-XL-TOPO	cloning vector	Invitrogen
pKSH827	cloning vector carrying <i>kshA3</i> ; KshARho-3-F + -R PCR (Expand polymerase) product (<i>kshA3</i>) ligated in pCR-XL-TOPO	This study
pKSH829	<i>kshA3</i> expression plasmid; 1,216 kb <i>NdeI/BamHI</i> fragment of pKSH 827 in pET15b <i>NdeI/BamHI</i>	This study
pA3rho2	co-expression construct <i>kshA3-kshB</i> ; 1,500 bp <i>HindIII/XbaI</i> fragment of pBrho5 in pKsh829 <i>HindIII/SpeI</i>	This study
pA4rho4	KshA4 and KshB co-expression vector	Petrusma <i>et al.</i> (2009) (24)
pKSH920	cloning vector carrying <i>kshA5</i> ; <i>kshA5</i> PCR product obtained using primers KshARho-5-F and KshRho-5-R ligated in pZero <i>EcoRV</i>	This study
pKSH921	<i>kshA5</i> expression plasmid; 1,227 bp <i>NdeI/BamHI</i> fragment of pKSH 920 ligated in pET15b <i>NdeI/BamHI</i>	This study
pA5rho2	co-expression construct <i>kshA5-kshB</i> ; 1,216 bp <i>HindIII/XbaI</i> fragment of pBrho5 ligated in pKsh921 <i>HindIII/SpeI</i>	This study